III. CONCLUSION

- 21. From the foregoing analysis, I find that Spezyme Ethyl contains DNA having the nucleotide sequence set forth in Exhibit 3. This DNA encodes a protein, which has the amino acid sequence that is also depicted in Exh. 3.
- 22. I declare under penalty of perjury pursuant to the laws of the United States of America that the foregoing statements are true and correct.

Respectfully submitted,

Dated: 19, May 2005

Steen Troels Jørgensen

Attachments:

Exhibit 1: Curriculum Vitae of Steen Troels Jørgensen

Exhibit 2: Spezyme Ethyl DNA Sequence

Exhibit 3: P.L. Jørgensen et al., "Cloning of a chromosomal α-amylase gene from Bacillus

stearothermophilus" FEMS Microbiology Letters (1991) 77:271-276.

EXHIBIT 1

Steen Troels Jørgensen Curriculum Vitae

Initials: STJq	Full name: Steen Troels Jørgensen				
Date of birth: 8. June 1958	Reports to (initials): MDRa				
Year of graduation: 1985	University/college: University of Copenhagen				
Degree: Cand. Scient	Joined NN in year: 1985	· · · · · · · · · · · · · · · · · · ·			

Job Experience:

Year:	Business unit/company:	Position:	Assignments/tasks:
1985	Bacterial Gene Technology, Novo Industri A/S.	Research Scientist	Gene technology as used in the construction of bacterial production strains for high-level expression of enzymes, and in the cloning and expression of novel enzyme coding genes.
1995	Bacterial Gene Technology, Enzyme Business, Novo Nordisk A/S	Principal Scientist (Science Manager)	As above.
2002	Bacterial Gene Technology, Novozymes A/S	Senior Science Manager	As above.

RESEARCH WORK & BIBLIOGRAPHY:

REGULAR PAPERS:

Bech, F. W., Jørgensen, S. T., Diderichsen, B., and Karlström, O. H. (1985). Sequence of the *relB* transcription unit from *Escherichia coli* and identification of the *relB* gene. The EMBO Journal, 4, 1059-1066.

Gerdes, K., Bech, F. W., Jørgensen, S. T., Løbner-Olesen, A., Rasmussen, P. B., Atlung, T., Boe, L., Karlstrøm, O., Molin, S., and von Meyenburg, K. (1986). Mechanism of postsegregational killing by the *hok* gene product of the *parB* system of plasmid R1 and its homology with the *relF* gene product of the *E. coli relB* operon. The EMBO Journal, 5, 2023-2029.

Jørgensen, S., Skov, K. W., Diderichsen, B. (1991). Cloning, sequence and expression of a lipase gene from *Pseudomonas cepacia*: Lipase production in heterologous hosts requires two *Pseudomonas* genes. J. Bacteriol., 173, 559-567.

Diderichsen, B., Poulsen, G. B., and Jørgensen, S. T. (1993). A useful cloning vector for *Bacillus subtilis*. Plasmid 30, 312-315.

Norman, B. E., Jørgensen, S. T. (1992). *Thermoanaerobacter* sp. CGTase: Its properties and application. Denpun Kagaku, 39, 99-106.

Hobson, A. H., Buckley, C. M., Aamand, J. L., Jørgensen, S. T., Diderichsen, B., and McConnell, D. J. (1993). Activation of a bacterial lipase by its chaperone. Proc. Natl. Acad. Sci. USA, 90, 5682-5686.

Kelly, A. P., Diderichsen, B., Jorgensen, S., McConnell, D. J. (1994). Molecular genetic analysis of the

Steen Troels Jørgensen Curriculum Vitae

pullulanase B gene of Bacillus acidopullulyticus. FEMS Microbiology Letters, 115, 97-106.

Aamand, J. L., Hobson, A. H., Buckley, C. M., Jørgensen, S. T., Diderichsen, B., and McConnell, D. J. (1994). Chaperone mediated activation in vivo of a *Pseudomonas cepacia* lipase. Mol. Gen. Genet. 245, 556-564.

Hobson, A. H., Buckley, C. M., Jørgensen, S. T., Diderichsen, B., and McConnell, D. J. (1995). Interaction of the *Pseudomonas cepacia* DSM3959 lipase with its chaperone LimA. J. Biochem., 118, 575-581.

Tangney, M., Jørgensen, P. L., Diderichsen, B, and Jørgensen, S. T. (1995). A new method for integration and stable DNA amplification in poorly transformable bacilli. FEMS Microbiology Letters 125, 107-114.

Pedersen, S., Jensen, B. F., Jørgensen, S. T. (1995). Enzymes from Genetically Modified Microorganisms. In "Genetically Modified Foods. Safety Aspects." Eds. Engel, K.-H., Takeoka, G. R., Teranishi, R. ACS Symposium Series 605, pp. 196-208.

Pedersen, S., Dijkhuizen, L., Dijkstra, B. W., Jensen, B. F., Jørgensen, S. T. (1995). A better enzyme for cyclodextrins. Chemtech vol. 25, no. 12, pp. 19-25.

Jørgensen, S., Vorgias, C. E., Antranikian, G. (1997). Cloning, sequencing, characterization, and expression of an extracellular α-amylase from the hyperthermophilic Archaeon *Pyrococcus furiosus* in *Escherichia coli and Bacillus subtilis*. J. Biol. Chem., 272, 16335-16342.

Jørgensen, S.T., Tangney, M., Starnes, R. L., Amemiya, K., Jørgensen, P. L. (1997). Cloning and nucleotide sequence of a thermostable cyclodextrin glycosyltransferase gene from *Thermoanaerobacter* sp. ATCC 53627 and its expression in *Escherichia coli*. Biotechnology Letters, 19, 1027-1031.

Jørgensen, S. T., Tangney, M., Jørgensen, P. L., Diderichsen, B. (1998). Integration and amplification of a cyclodextrin glycosyltransferase gene from *Thermoanaerobacter* sp. ATCC53627 on the *Bacillus subtilis* chromosome. Biotechnology Techniques, 12, 15-19.

Krogh, S., Jørgensen, S. T., Devine, K. M. (1998). Lysis genes of the *Bacillus subtilis* defective prophage PBSX. J. Bacteriol. 180, 2110-2117.

Jørgensen, P. L., Tangney, M., Pedersen, P. E., Hastrup, S., Diderichsen, B., Jørgensen, S. T. (2000). Cloning and sequencing of an alkaline protease gene from *Bacillus lentus* and amplification of the gene on the *B. lentus* chromosome by an improved technique. Appl. Environ. Microbiol. 66, 825-827.

Stephenson, K., Jensen, C. L., Jørgensen, S. T., Lakey, J. H., Harwood, C. R. (2000). The influence of secretory-protein charge on late stages of secretion from the gram-positive bacterium *Bacillus subtilis*. Biochem. J., 350, 31-39.

Jensen, C. L., Stephenson, K., Jørgensen, S. T., Harwood, C. (2000). Cell-associated degradation affects the yield of secreted engineered and heterologous proteins in the *Bacillus subtilis* expression system. Microbiology, 146, 2583-2594.

Outtrup, H., and Jørgensen, S. T. (2002). The Importance of Bacillus species in the Production of Industrial Enzymes. Chapter 14, pp. 206-218, in Berkeley, R., Heyndrickx, M. Logan, N. and De Vos, P (eds.) "Applications and Systematics of *Bacillus* and Relatives". 2002, Blackwell Science Ltd.

Stephenson, K., Jensen, C. L., Jørgensen, S. T., Harwood, C. R. (2002). Simultaneous inactivation of the wprA and dltB genes of Bacillus subtilis reduces the yield of α -amylase. Letters in Applied Microbiology, 34, 394-397.

Craynest, M., Jørgensen, S., Sarvas, M., Kontinen, V. P. (2003). Enhanced secretion of heterologous cyclodextrin glycosyltransferase by a mutant of *Bacillus licheniformis* defective in the D-alanylation of

Steen Troels Jørgensen Curriculum Vitae

teichoic acids. Letters in Applied Microbiology, 37, 75-80.

POSTERS AND PRESENTATIONS:

Steen Jørgensen, Kirsten W. Skov, and Børge Diderichsen (1990). A novel phenomenon involved in heterologous expression of a bacterial lipase gene: requirement for an additional gene from the donor organism. Oral presentation July 10, 1990, at the 5th European Congress on Biotechnology, Copenhagen, by S. Jørgensen. Abstract book p. 60.

Norman, B. E., and Jørgensen, S. T. (1992). Thermoanaerobacter sp. CGTase: Its properties and application. Denpun Kagaku, 39, 99-106. (Proceedings of the amylase symposium, 1991. Oral presentation by B. E. Norman).

Starnes, R. L., Amemiya, K., Jørgensen, P. L., and Jørgensen, S. (1992). Cloning and expression of the Thermoanaerobacter sp. Cyclodextrin glycosyl transferase gene in Escherichia coli and Bacillus subtilis. Oral presentation at Sixth International Cyclodextrin Symposium, April 21-24, 1992, Chicago, USA, by R. L. Starnes.

J. L. Aamand, C. M. Buckley, A. H. Hobson, S. Joergensen, B. Diderichsen, and D. J. McConnell (1992). Expression of a Pseudomonas cepacia lipase in E. coli.

Poster presented at the Protein Engineering Workshop in Korpilampi, Finland, 12-17 August 1992, by J. L. Aamand.

Audrey H. Hobson, Catherine M. Buckley, Steen Jørgensen, Børge Diderichsen, and David J. McConnell (1992). Conformational activation of a bacterial lipase.

Poster presented at the conference "Understanding Protein Folding", Novum Research Centre, Stockholm, 24-28 August 1992, by A. H. Hobson.

Jørgensen, S. T., Jørgensen, P. L., Diderichsen, B. (1993), Genetic stabilization of recombinant industrial Bacillus species for large scale fermentation of industrial enzymes. Oral presentation at 7th International Bacillus Conference, Institut Pasteur, Paris, on July 20, 1993, by Børge Diderichsen.

Antranikian, G., Rüdiger, A., Canganella, F., Jørgensen, S. (1994). Enzymes from hyperthermophiles; especially starch degrading enzymes.

Oral presentation at 7th International Symposium on the Genetics of Industrial Microorganisms, June 26-July 1, 1994, Montreal, Canada, by G. Antranikian.

Pedersen, S., Jensen, B. F., Jørgensen, S. T. (1994). Enzymes from genetically modified microorgan-

Oral presentation at the 208th ACS National Meeting, Washington DC, August 22, 1994, by Sven Pedersen.

Jensen, C. L., Stephenson, K., Jørgensen, S. T., Harwood, C. R. (1995). Construction of hybrid αamylase genes for investigation of the interaction between secretory proteins and the Bacillus subtilis cell wall.

Poster presented at the 8th international conference on Bacilli, July 8-12, 1995, Stanford, California.

Jørgensen, S. T. (1996). Prospects and problems in large scale production with recombinant Bacillus. Oral presentation at the Bioscience Days, 14. September 1996, Helsinki, Finland.

Jorgensen, S. T., Jensen, C. L., Kristensen, T., Nielsen, B. R., Stephenson, K., Harwood, C. R. (1997). Construction and use of hybrid lpha-amylase genes for investigation of protein secretion from Bacillus subtilis. Poster presented at European Cell Factory Conference, Lund 27-30 April 1997.

Jensen, C. L., Stephenson, K., Harwood, C. R., Jørgensen, S. T. (1997). Protein secretion from Bacil-

Steen Troels Jørgensen Curriculum Vitae

lus subtilis studied using chimeric α -amylases with changed net charge. Oral presentation at 9^{th} International Conference on Bacilli, Lausanne, Switzerland, 15-19 July 1997, by C. L. Jensen.

Jørgensen, S. T. (1997). A general, transposon-based method for the construction of selection marker free *Bācillus* strains containing multiple, chromosomally integrated heterologous genes. Oral presentation at 9th International Conference on Bacilli, Lausanne, Switzerland, 15-19 July 1997, by S. T. Jørgensen.

Stephenson, K., Jensen, C. L., Jørgensen, S. T., Petit-Glatron, M.-F., Chambert, R., Harwood, C. R. (1997). Factors affecting late stages in the secretion of α -amylases from Bacillus subtilis. Poster presented at 9th International Conference on Bacilli, Lausanne, Switzerland, 15-19 July 1997.

Stephenson, K., Jensen, C. L., Carter, N. M., Jørgensen, S. T., Petit-Glatron, M.-F., Chambert, R., Harwood, C. R. (1998). Late stages in the secretion of proteins from *Bacillus subtilis*. Oral presentation at the International Conference on Bacilli, Japan, July 13, 1998, Osaka, by. Stephenson, K.

Rasmussen, M. D., Jørgensen, S. T., Jensen, C. L., Jørgensen, P. L. (1999). Industrial approach for cloning and expression optimization of α -amylases. Oral presentation at Society for Industrial Microbiology 50th Anniversary Annual Meeting, August 1-5, 1999, Crystal Gateway Marriott, Arlington, VA., by M. D. Rasmussen.

PATENT APPLICATIONS/PATENTS:

Jørgensen, S. T. (1991). Dna encoding a lipase and a lipase modulator. PCT publication WO 91/00908. European Patent Specification O 487 541 B1.

Jørgensen, S. T., Jørgensen, P. L., and Diderichsen, B. K. (1991). Stable integration of DNA in bacterial genomes. PCT publication WO 91/09129. US Patent 5,695,976. European Patent Specification 0 506 780 B1.

Antranikian, G., Jørgensen, P. L., Wümpelmann, M., Jørgensen, S. T. (1992). Novel thermostable pullulanases. PCT publication WO 92/02614.

Jørgensen, S. T., Diderichsen, B. K., Buckley, C. M., Hobson, A., Mcconnell, D. J. (1993). A process for the preparation of lipase. PCT publication WO 93/13200. US Patent 5,681,715.

Jørgensen, S. T., and Jørgensen, P. L. (1993). A process for expressing genes in *Bacillus licheniformis*. PCT publication WO 93/10248.

Jørgensen, S. T, and Diderichsen, B. K. (1993). A Bacillus promoter derived from a variant of a Bacillus licheniformis α -amylase promoter. PCT publication WO 93/10249. US Patent 5,698,415.

Jørgensen, S. T. (1994). DNA amplification. PCT publication WO 94/14968. US Patents 5,733,753 and 5,925,544.

Jørgensen, S. T. (1994). An amylolytic enzyme. PCT publication WO 94/19454.

Jørgensen, S. T. (1996). DNA integration by transposition. PCT publication WO 96/23073. US Patent 5,882,888.

Tangney, M., Hansen, C., Pedersen. P. E., Jørgensen, P. L., Jørgensen, S. T. (1996). Introduction of DNA into Bacillus strains by conjugation. PCT publication WO 96/29418. US Patents 5,843,720 and 6,066,473.

Fuglsang, C. C., Okkels, J. S., Petersen, D. A., Patkar, S., A., Thellersen, M., Vind, J., Halkier, T., Jørgensen, S. T. (1997). A modified enzyme with lipolytic activity. PCT publication WO 97/04079.

Steen Troels Jørgensen Curriculum Vitae

US Patent 6,495,357.

Adams, L. F., Widner, W. R., Diderichsen, B. K., Thomas, M. D., Jorgensen, S. T., Sloma, A. P., Jorgensen, P. L. (1997). Production of Bacillus thuringiensis integrants. PCT publication WO 97/27305. US patent 5.955,367.

Harwood, C.R., Stephenson, K., Jorgensen, S., Hansen, C., Kristensen, T. (1998). Improved prokary-otic expression of protein. PCT publication WO 98/49328.

Jørgensen, S. T., Pedersen, K. B. (1999). Protein producing cell containing multiple copies of a desired gene and a screenable marker but no selection marker. PCT publication WO 99/01562.

Jørgensen, S. T., Frandsen, T. P., Andersen, C. (1999). Expression of starch-binding domain (SBD). PCT publication WO 99/15636.

Jørgensen, S. T. (1999). A prokaryotic cell comprising two copies of a gene transcribed in different directions. PCT publication WO 99/41358. US patent 6,100,063.

Joergensen, S. T., Christensen, C. L., Kristensen, T. (2000). A *Bacillus* protein production cell. PCT publication WO 00/63346.

Jørgensen, S. T., Andersen, J. T., Rasmussen, M. D., Olsen, C. (2002). Method for stable chromosomal multi-copy integration of genes. PCT publication WO 02/00907

Jørgensen, S. T., Andersen, J. T., Banke, N., Nielsen, P. (2003). Eubacterial RNA-polymerase mutants with altered product production. PCT publication WO 03/055996.

Andersen, J. T., Jørgensen, S. T., Rasmussen, M. D., Olsen, P. B. (2003). Improved Bacillus host cell. PCT publication WO 03/093453.

Andersen, J. T., Jørgensen, S. T., Rasmussen, M. D., Olsen, P. B. (2003). Improved Bacillus host cell. PCT publication WO 03/087148.

Andersen, J. T., Jørgensen, S. T., Rasmussen, M. D., Olsen, P. B. (2003). Improved Bacillus host cell. PCT publication WO 03/087149.

Andersen, J. T., Jørgensen, S. T., Rasmussen, M. D., Olsen, P. B. (2003). Improved Bacillus host cell. PCT publication WO 03/087142.

EXHIBIT 2

FEMS Microbiology Letters 77 (1991) 271-276
©1991 Federation of European Microbiological Societies 0378-1097/91/\$03.50

ADONIS 037810979100051S

271

FEMSLE 04275

Cloning of a chromosomal α -amylase gene from *Bacillus stearothermophilus*

Per Linå Jørgensen, Gitte Bak Poulsen and Børge Diderichsen

Novo Nordisk, Bagsværd, Denmark

Received 22 August 1990 Accepted 18 September 1990

Key words: Native plasmids; Heat stable amylases; Nucleotide sequence; Evolution; Cloning in Bacillus subtilis

1. SUMMARY

We have cloned and sequenced a gene for a heat-stable α -amylase from a natural isolate of *Bacillus stearothermophilus*. Previously, it had been shown that *B. stearothermophilus* amylase genes may be harboured on indigenous plasmids. We have found that our isolate harbours the amylase gene only on the chromosome and not on its indigenous plasmid.

2. INTRODUCTION

Amylases are exoenzymes, which are widely distributed in nature. All amylases, which have been sequenced so far, share significant homology suggesting an evolutionary relationship. The observation that an α -amylase gene from B. stearothermophilus is harboured on a natural plasmid [1] supports the notion that gene transfer can take place between related strains. In this paper we show that homologous amylase genes

can reside either on a native plasmid or on the chromosome of different strains of *B. stearother-mophilus*.

3. MATERIALS AND METHODS

3.1. Bacterial strains and plasmids

Escherichia coli K-12 MT102 is (araD139, Δ (ara-leu)7697, lacX74, galU, galK, rpsL, r⁻ m⁺). Bacillus subtilis DN1885 (amyE, amyR2) is a derivative of B. subtilis 168 strain RUB200 [2]. B. stearothermophilus DN1792 and DN1793 were isolated from a household hot water system and classified by Deutsche Sammlung für Mikroorganismen. ATCC 31783 is a B. stearothermophilus from the American Type Culture Collection.

Plasmid pUN121 has been described [3]. Plasmid pDN2360 was constructed by ligation of PstI fragments from pUN121 and plasmid pDN1380 [4] (Fig. 1). Plasmid pBB37 (Fig. 1) was constructed by inserting a ClaI fragment harbouring the gene for an α -amylase from B. amyloliquefaciens [5] into the ClaI site of plasmid pDN1620, which was derived from pDN1380 by insertion of an oligonucleotide linker in the EcoRI site.

Correspondence to: B. Diderichsen, Novo Nordisk, DK-2880 Bagsvaerd, Denmark.




Fig. 1. Restriction maps of plasmids. pDN2360 replicates in both E. coli and in B. subtilis. pBB37 and pPL1059 replicate in B. subtilis. cI is the coliphage λ repressor gene. P_r is a promoter from λ. P_m is the promoter of the maltogenic amylase gene amyM [4]. cat is the chloramphenical resistance conferring gene from plasmid pC194 of Staphylococcus aureus. amyQ is the amylase gene from B. amyloliquefaciens [5]. P_q is the promoter from the amyQ gene. amyS is the amylase gene from B. stearothermophilus described in this work.

3.2. Transformation and plasmid preparation

Competent cells, transformation, and preparation of recombinant plasmids were as described [4]. Transformants of E. coli were selected with 7 μ g/ml tetracycline. Transformants of B. subtilis were selected with 6 μ g/ml chloramphenicol.

p	25	- K	D.					
1300 1310 1320 1330 1340 1350 1350 1360 AACCOTYGGCTTAACGCCTATAATACGAAAAAGAAAGAAAGAAAGAA	1170 1180 1190 1400 1410 1420 1430 1430 1430 1430 1430 1430 1430 143	1510 	1580 	1590 1600 1610 1620 1630 1640 1650 AACACCTCGAAAACACTTTTTCTTACCTTACCGAATGATGATCATCTTACCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCTTACATCAT	~	1730 1740 1750 1760 1770 1770 1790 1790 COGCCGARCCAACCCAACCCAACCCCAACCCAACCCAACC		
1360° 	1430 1 acacaca R R 1		2116	1650 	1720 1720 1721 1721 1721	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$		
100	1420 1430 1 	1500 CAAGGGG	20 XXX	. 25×	·	1790 - Trodra		
1350 TCHITAT	TCGC	3. 8 3. 8	1570 1 NGCATGERAC	1630 1640 	9 8	3700		
3-62	1429 CHCK	1490 GGACAN	Z,	, YOU'	1710 1710 XXCG	1780 1 1 1 1 1 1 1 1 1		
200	G.	21 DE 2	1560 	8 C	. Se ×	NCCG		
1340 GATACCCG	NTCC	2001	33,056	1630 	1700 1 7 7 V P	1760 1770 1 1 1 170 170 170 170 170		
GGAT	1400 1410 CTGAAAAGCAAAATCGATH L K S K I D	1480 1 37CK	g 95%	CGGN	× X3667 J	OLTI L R V		
1330 CAGGAA O E	3,×	CGAC	1550 1 CATGGGCC0 D G P	1620 1 1 0653300	0 E >	, F		
1330 - 13665/66	MAAGE x	1470 VTCACTC		NCCOCK	1690 - -	1760 Janaa 1760 1760		
TOTAL PER	r tag	75 P	1540 HATCACCG I T	1610 NCCTTA D L	5005	1,000		
1320 Trecta	1390 CCTTCG P S	E a	1530 154 Creececacrearc L. A. A. L. I.	1600 1610 	168b 	1750 		
T.	139 NTYCCT 1 P	1460 	1530 	Y X	Arco N	1750 		
1310 	CANCO N	· KC	1 2 2	1600 	1670 Julyerca K	ACCC T		
T K	1380 - XXTAX	1450 - ACGCA) T 0	900	3×	AL AND	1740 CACA	1810 	
Tree	CCAC	1450 1460 1470 1480 1490 262ACACACAGATACACCACCACAGATACACCACCACATACAT	1520 I NACCAGUATCCC X P G 5	1590 	XXII E	1740 E I T	118	
AACCGTTG	1370 CATE G I	TTAC	ACC.N(AACACGCT	1660 - 	1730 CTCGGC R	TEAT	•
	790 1. Proc 9	800 810 920 830 840 850 860 1450 1460 1470 1480 1490 1500 1510 1510 1510 1510 1510 1510 15	<	₹ o 2		2 E	1160 1170 1180 1190 1200 1210 1220 1810 1810 CCMARCARANTAGE CTARATAGE CTARATAGA CTARATAGE CTARATAGA CTARAT	A.
650 660 670 680 690 700 710 CGGCCACTAYTCAAATCAAAATTTAATTTTTCCGGGGGGGCAAAACCAACTAAA S G T Y Q I Q A W T K T D F P G R G H T Y S S F	730 740 750 760 770 780 780 790 770 780 790 780 780 780 780 780 780 780 780 780 78	640 850 860 GGAACTATGATGCCGGAAACTATGACTATGCCTACTATGACTATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGGAAACTATGATGCCGAAACTATGATGATGCCGAAACTATGATGCCGAAACTATGATGCCGAAACTATGATGCCGAAACTATGATGATGCCGAAACTATGATGATGATGATGATGATGATGATGATGATGATGATG	910 920 930 	940 950 960 970 980 990 1000 ACATANATOGOTICGGCTTANGCCGCCANACANTANGCTCANTOGATTTTTTCCGANTGATTGATT N I D OF R L D A V K H I K F S F F P D W L S Y	1910 1920 1939 1940 1950 1960 1970 ACTICACACCACCACTATTACCGCGGGGAAAAATTGGACAAACACACAC	090 1100 1110 1120 1130 1140 1150 1150 1150 1150 1150 1150 115	1220 - CATTG	SK 5 G G A F D H R T L H T N T L H K D G P T L 1230 1240 1250 1250 1250 1250 1290 CCGTCACTTCGTTCATATATATATATACAACCCGGCAAGCCGTCAAGCGTCAAGCGTCAAGCGTCAAGCGTCAAGCGTAAGCGTAACAGCGTAAGCGTAAGCAACAGCAAGCA
710 FACTCCA(Y S :	TXCA.	XIGT.	MCN.	1000 1000 11000 11000	סר האמדי	ENTAC Y 1	CCGAC	P T 1290 CCATGO
ACCTAC T Y	780 CATT	850 SCTTA	v v	7.170 7.	CAACAAG	MATTY	1210 GATCAN	ဝ ၌
NACAC H T	S B	8-K	920 	990 ICCTGA	ACAT	NCX.	27 - 89	x D 1280 1280 1280 1280
700 - - 0 H	AWITEN	, v	X,X	E_	1060 - TATE!	1130 Trickch	XTGX.	r SCAT
) 100000 100000	EGAAA.	BANKE N	910 1 196660	TE T	GAGC S	11.67	1200 1 27 27 27	T L 1270 - - - -
069	AGCC(50 ×	,	980 - - - - - - - -	1050 GANTATTGG/ E Y W	, ECCC	ATAC	z 27-67
69 17777771 8 8	760 1 1 3ACGAAAG D Z S	830 CGGAAU T E	. axx	55×	T TOWN	1120 	1190 I VTGACCA	မ ဦ
o y	regan V D	ACAC D	900 - 	960 970 94 	40 Greasac	TOTE	I. M.	1260 1260 1260 1260
680 XXXTT	750 TTTGATT	O V	ACCG T	×	1040 - Accord T	1110 	Accr	r Sy
EACG	GGCGTT G V	820 1 1 1 1 1 1 1 1	890 1 1 1 1 1 V	o carc	1030 10 	DATG	1180 00000	И В 1250 ј ј 137САСС
670 - 	740 i Treaced F D g	D.T.T.C	NAGE V	960 	1030 - - - - - -	1100 1 ACAGACGGAACG T D G T	KTAT	2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
OCAN	4 -FF	810 7 77000	880 XTCCCC	i i	, XXG	1100 1 100 100 100 100	0711 1 1 1 1 1 1 1 1 1 1	* 9 XX
660 NTCANATC Y Q I	D PACCAT	7 Y	MTCX.	950 	700CU	35	2000	1240 1240 W. G.
- 57.×	730 : (CTGGGT (W R	aon Coccx	X 756	0007	1020 CAGACT	1090 	1160 	LICCI C
SCACC G T	א א	CATC	B70 TGXT	940 Grann	TTCT		11-27	1230 1230 124CC
650 CGGG 5	AGTO K	9000	ACCT D L	ACAT X	1010 - TGCGT N	XTTAC N Y	3	2 CCG7 X
2-E	9				0 5	ų		
C _FO	- E	210 TTGCT 1. L	STGCA M D	TATC	NCGG Y	S00 NAAGCT A A	570 GACCA	o o o
Je je je	[TGA]	. Y	28. - 28. 28. 28.	350 2AAC1	\GGG7	ACO.	Parte	F D
04 100	130 - -	150 170 180 190 200 210 DOCHAGETTCACCEARCATCADAMGANGATCACTCACCTTRITTCAC HR I I R K G W W F L L A F L L	230 240 250 260 270 280 100 100 100 100 100 100 100 100 100 1	100 110 120 130 140 150 SECTION SECURITY SECURIT	420 J ACGTAC D V	490 	o rcore	A CONTRACTOR
CGNG	1 TOTA	C. C. L.	270 	G. N.G.	AGCG S	XXX	560 - -	630 630 70 70 70 70 70 70 70 70 70 70 70 70 70
Se - Se Corri	CGAA	GITC	Ę	CAAT	410 	o Sch	2000	۲) عرقن
TAA.	120 - 	190 FGGAT	So se	TGGC	e s	480 	550 1 STCTA	620 620 1 1 1 1 1 1 1 1 1 1 1
) IGACG	, (S.	VOGW1	260 1 1 7 7 8	*8-33×	400 H CAAGGAA K	TACCG	CAAG	o co
TAXEA TA	110 - 1200 1300 1300 1300 1300 1300 1300 1300	180 CGMAAA R K	ומפנד ר	T T	¥-8×	4.70 2.000 2.000 2.000	540 	E CGN
VEEN	13.00	i – pr	250 1000 1000 1000	320 	E. C	3,5	s – s	610 610 610 610 610 610
8- 8	ATT	EMIC.	4 122201	E - G	3300 - 10000 -	160 FCAAT	າວວອນ	د م م
EXXX	100 - CTTAX	170 170 18 18	240 CGACX	ATGG	TGCC	A-my	530	A H 600 1 VIGGO
TTAC	7700	irric.	ACAG T	310 - 96AT007	380 1 1 1	5000	5005	A PER
ACGA	90 12CGT	150 TAACG E T	A CCC	2000	E Line	450 CCTCC	520 TTCM	CACG
CON	rackA	moc:	230 F C	300 - - - -	, recec	XTOX Y D	CCAI	590 C
OL LEATE	80 	CATT	icnor 1	Taga	370 	440) CTTGI	D TCAAG	ه التوديد
10 20 39 40 50 70 70 10 10 10 10 10 10 10 10 10 10 10 10 10	BO 90 100 110 120 130 140 140 150 150 150 150 150 150 150 150 150 15	150 160 170 180 190 200 210 GGAAGGGCATTTGGAAAAGAAGAATGATCCTGCTGTTTTGCTCTATTGAAAAGAATGATCTTGTGCTTTTTGCTCTATTGCTATTGCTCTATTGCTCTATTGCTCTATTGCTCTATTGCTCTATTGCTCTATTGCTCTATTGCTATTGCTCTATTG	120 230 240 250 260 270 280 270 CHOROCOCCANGCOCA	290 100 110 120 130 140 350 ATTTACANGGARACCHAGGARACCANAGACCARANCATANCCA Y F E W Y L F P D G T L W T K V A M E A M H L S	370 380 390 400 410 420 430 accidentes accid	440 450 460 470 480 490 500 NINGENTATION CONTROL OF THE STATE OF THE	NO STO STO STO STO STO STO STO STO STO ST	OYLOAIQAAHAAGHOOFE OO SOO 600 600 600 600 600 600 600 600 600 6
XX.	TAC	CGA	P. F	u-f'	ស ស	EXT.	125	٥ ٢٤

Fig. 2. DNA sequence of the anyS gene. Promoter regions -35 and -10 and a ribosome binding site have been underlined starting at bp 11, 35 and 140, respectively.

An arrow shows the processing site after the 34 amino acid signal peptide.

273

274

Large plasmids were prepared by alkaline lysis according to ref. 6 with the exception that 50 mM glucose was added to the suspension buffer and that the samples were centrifuged for only 10 min before phenol extraction. Chromosomal DNA was prepared by phenol extraction [7].

3.3. DNA sequencing

DNA sequencing was performed directly on doublestranded plasmid templates using the SequenaseTM kit from United States Biochemical Corporation.

3.4. Southern analyses

Chromosomal and plasmid DNA was digested with restriction enzymes, fractionated on 0.7% agarose gels and blotted onto nitrocellulose filters. pPL1022 was labelled by nick translation using [³²P]dCTP (Amersham). The hybridization was according to Southern [8].

4. RESULTS

4.1. Cloning in E. coli

Genomic DNA from strain DN1792 was partially digested with Sau3A and fragments, of 3-10 kb were ligated with the BclI digested selection vector pUN121. E. coli MT102 was transformed with the ligation mixture selecting for tetracycline resistance on agar plates containing starch. Of 10000 transformants, two had a clear amylase positive phenotype. One of these contained a plasmid, pPL1011, with an insert of 8 kb. pPL1011 was partially digested with Sau3A and ligated with selection vector pDN2360 digested with Bcl I. Tetracycline resistant transformants of E. coli MT102 were screened for amylase expression and one positive transformant harbouring pPL1022 with an insert of 2.2 kb was kept for further study. The structural gene for the B. stearothermophilus amylase was called amyS.

4.2. Cloning in B. subtilis

A 2.0 kb *DraI-HindIII* fragment of pPL1022 was ligated with the 3.3 kb fragment of *B. subtilis* plasmid pBB37 and transformed into *B. subtilis* DN1885 selecting for chloramphenicol resistance

and screening for amylase activity. A positive transformant harboured the 5.3 kb plasmid pPL1059 (Fig. 1).

4.3. DNA sequence analysis

The sequence of the amyS gene of pPL1022 is shown in Fig. 2. Of the 1814 bp of this sequence, 33 bp (1.8%) differed from the sequence of Nakajima et al. [9] and 40 bp (2.2%) from the sequence of Gray et al. [10]. Comparing the amino acid sequence of the mature enzyme and the signal peptide with the above sequences showed 12 (2.2%) and 9 (1.6%) differences, respectively. Between the sequences of Nakajima et al. [9] and Gray et al. [10], there are 13 (2.4%) amino acid differences.

At the amino acid level, one of the three sequences differs from the two others at 17 positions of which 4 (11.8%) are in the signal peptide of 34 amino acids and the remaining 13 (2.5%) in the mature part of 515 amino acids.


4.4. Southern blotting analysis

To test the localization in *B. stearothermophilus* of the cloned amylase gene, plasmids were prepared from the *B. stearothermophilus* DN1792, DN1793 and ATCC 31783 [11]. DN1792 and DN1793 contained each one plasmid of 25 kb and ATCC 31783 contained three plasmids of 12, 15 and 26 kb as reported by Mielenz [1] (Fig. 3A). Using the cloned amylase gene as probe, hybridization was only seen with the 26 kb plasmid of ATCC 31783. The probe, however, clearly hybridized with the chromosomal DNA of DN1792 and DN1793 (Fig. 3B). Thus, *amyS* is located on the chromosome of DN1792 and DN1793 and on a native plasmid in ATCC 31783.

5. DISCUSSION

The amyS gene of a natural isolate of B. stearothermophilus has been cloned, sequenced and compared with genes from other B. stearothermophilus strains. The sequences of three independently isolated amyS genes (refs. 9, 10, and this work) show some variations in particular in the signal peptide regions.

A1 2 3 4 5 6 7 8 9 10 B1 2 3 4 5 6 7 8 9






Fig. 3. Southern blotting analysis of *B. stearothermophilus* plasmids and chromosomal DNA. (A) 0.7% agarose gels stained by ethidium bromide. (B) The same gels as in A but hybridized with nicktranslated pPL1022. Lanes 1 and 4, plasmid from DN1792; lanes 2 and 5, plasmid from DN1793; lanes 3 and 6, plasmid from ATCC 31783; lanes 7–9, chromosomal DNA from DN1792, DN1793 and ATCC31783, respectively; lane 10, λ DNA digested by *Bst* E2. Lanes 1–3, uncut plasmids; Lanes 4–6, plasmids cut by *HindIII*; Lanes 7–9, chromosomal DNA cut by *Eco* RI.

Satoh et al. [12] have cloned two closely related amylase genes from what appear to be two phylogenetically very distant B. stearothermophilus strains. Nucleotide sequence analyses show that the regions flanking the amylase genes are completely different. Sen and Ariel [13] report on the location of amylase genes in two B. stearothermophilus strains. In one strain, the amylase gene is present both on two different indigenous plasmids and in multiple copies on the chromosome. In the other strain, multiple copies were found on the chromosome but no indigenous plasmid could be detected. We speculate that the multiple chromosomal genes in these two strains may be the result of plasmid integration. In this paper, it is shown that two other B. stearothermophilus strains harbour the amylase gene on the chromosome only, despite the presence of indigenous plasmids. It is thus likely that there is a mechanism in B. stearothermophilus which allows for migration of amylase genes from chromosome to plasmid or vice versa.

REFERENCES

- [1] Mielenz, J.R. (1983) Proc. Natl. Acad. Sci. U.S.A. 80, 5975–5979.
- [2] Yoneda, Y., Graham, S. and Young, F.E. (1979) Biophys. Biochem. Res. Commun. 91, 1556-1564.
- [3] Nilsson, B., Uhlén, M., Josephson, S., Gatenbeck, S. and Philipson, L. (1983) Nucleic Acids Res. 11, 8019–8029.
- [4] Diderichsen, B. and Christiansen, L. (1988) FEMS Microbiol. Lett. 56, 53-60.
- [5] Takkinen, K., Petterson, R.F., Kalkkinen, N., Palva, I., Söderlund, H. and Kääriäinen, L. (1983) J. Biol. Chem. 258, 1007-1013.
- [6] Maniatis, T., E.F. Fritsch, E.F. and Sambrook, J. (1982) Molecular Cloning, Cold Spring Harbor Laboratory, N.Y.
- [7] Yasbin, R.E., Wilson, G.A. and Young, F.E. (1975), J. Bacteriol. 121, 296-304.
- [8] Southern, E.M. (1975) J. Mol. Biol. 98, 503-517.
- [9] Nakajima, R., Imanaka, T. and Aiba, S. (1985) J. Bacteriol. 163, 401–406.
- [10] Gray, G.L., Mainzer, S.E., Rey, M.W., Lamsa, M.L., Kindle, K.L., Carmona, C. and Requadt, C. (1986) J. Bacteriol. 166, 635-643.
- [11] Mielenz, J.R. and Mickel, S. (1985) U.S. Patent 4, 493, 893.
- [12] Satoh, H., Nishida, H. and Isono, K. (1988) J. Bacteriol. 170, 1034-1040.
- [13] Sen, S. and Oriel, P. (1989) Gene 76, 137-144.